

## **Combined Manual Therapy Techniques for the Treatment of Women With Infertility: A Case Series**

Mary Ellen Kramp, DPT, CLT-LANA

Current treatment options for infertility, including hormone therapy, intrauterine insemination, and in vitro fertilization, tend to be expensive, are not necessarily covered by insurance, and carry different levels of short-term and long-term health risks. Many of the issues that contribute to infertility can be traced to scar tissue, fascial restriction, and lymphatic congestion in the pelvic region. Manual therapy techniques exist to release fascial restrictions, to mobilize tight ligaments, and to drain congested lymphatics, all of which can be applied to the reproductive system. In this case series, 10 infertile women were treated with 1 to 6 sessions of manual therapy applied to the pelvic region. Techniques included muscle energy, lymphatic drainage, and visceral manipulation. Six of the 10 women conceived within 3 months of the last treatment session, and all 6 of those women delivered at full term.

*J Am Osteopath Assoc.* 2012;112(10):680-685

Women in their childbearing years who have not been able to conceive after 1 year of unprotected intercourse are considered infertile according to the Centers for Disease Control and Prevention (CDC).<sup>1</sup> The CDC reports that 6.7 million women aged 15 to 44 years have impaired fecundity, and 1.5 million married women are infertile.<sup>1</sup> Worldwide, 15% of couples are reported as infertile.<sup>2</sup> In general, 27% of cases of infertility are caused by ovulation disorders; 25%, male factors; 22%, tubal disorders; 17%, unexplained factors; 5%, endometriosis; and 4%, other factors.<sup>3,4</sup> According to *Williams Gynecology*,<sup>4</sup> women who have not previously been able to conceive are considered to have primary infertility. Those who have previously conceived, whether or not the pregnancy was successful, are considered to have secondary infertility.

Average fertile women (aged 22-40 years) who have coitus in the week prior to ovulation have a 20% chance of developing a clinical pregnancy during each ovulatory cycle. Fifty-seven percent of fertile couples will conceive in the first 3 months, 72% in 6 months, and 85% in 1 year.<sup>4</sup>

Women whose infertility is unexplained have monthly fecundity rates of 10% to 15% with hormone therapy and intrauterine insemination (IUI).<sup>3</sup> In Canada, Collins et al<sup>5</sup> found that pregnancy rates for 873 infertile couples without any treatment were 35% after 3 years and 45% after 7 years. In the Netherlands, van der Steeg<sup>6</sup> found that, overall, untreated infertile couples were able to achieve spontaneous pregnancy 29.5% of the time within 12 months. Katz<sup>7</sup> reported that the incidence of infertility increases steadily in women after the age of 30.

Current options for treatment are dependent on the cause of infertility. Treatment options include fertility drugs, IUI, and in vitro fertilization (IVF). However, these treatments come with several considerations. First, procedures such as IVF are invasive and carry risk of infection. Second, the treatments can be expensive; the median cost for IVF is \$24,373.<sup>4</sup> It is often not covered by insurance and frequently needs to be repeated.<sup>4</sup> Success rates are approximately 38%,<sup>3</sup> and the successes have a high rate of multiple births, which places increased risks on the mother and the

fetuses.<sup>7</sup> Even though fewer embryos are being implanted now than a decade ago, multiple pregnancy remains the single highest risk of IVF.<sup>8</sup>

According to 2010 data on assisted reproductive technology, women attempting to conceive by means of IVF are doing so because of tubal factors (7%), ovulatory dysfunction (7%), diminished ovarian reserve (15%), endometriosis (4%), uterine factors (1%), male factors (17%), other factors (7%), and unknown factors (12%).<sup>8</sup> In couples undergoing IVF, 11% reported multiple factors in women and 18% reported multiple factors in both men and women.<sup>8</sup> In 2010, the national percentage of IVF cycles using fresh embryos from nondonor eggs that resulted in live births ranged from 5.0% to 41.5%, depending on age group.<sup>9</sup> Of cycles that resulted in conception, 56.6% resulted in a singleton pregnancy, 24.9% resulted in a multiple fetus pregnancy, and 16.4% ended in miscarriage, 0.9% induced abortion, 0.7% stillbirth, and 0.4% unknown outcome.<sup>10</sup> Pregnancy rates after IUI, according to Katz,<sup>4</sup> are 25% to 35%.

### ***Physiologic Factors in Infertility***

Many of the issues that cause a woman to have difficulty with conception can be traced to scar tissue, fascial restriction, and congested lymphatics.<sup>11,12</sup> Manual medicine has been used to manage these specific problems, but, to my knowledge, it has not been used to manage functional infertility. Part of the basic foundation of osteopathic medicine is that manual mobilization has an impact at the cellular level:

*The cells, which make up our body, have an internal environment also. The fluid matrix of it must be free of 'pollution.' Waste products of tissue metabolism must be constantly carried away by the veins and lymphatics. The health and life of cells and therefore the whole body depend on it.<sup>13</sup>*

To understand the concept of manual therapy as a treatment for patients with infertility, one must consider the reproductive environment at the cellular level and the reproductive anatomy at the tissue level. The arterial, venous, and lymphatic circulations are interrelated. The arteries transport the blood to the cells, then the venous system returns 90% of what was transported back to the

heart.<sup>12</sup> The lymphatic system will pick up particles too large to transport across the venous capillary membranes and interstitial fluid. Once interstitial fluid enters the lymph capillaries, it is called lymph.<sup>11</sup> The lymphatic system transports the remainder of this fluid (10% of the original fluid transported by the arteries) to lymph nodes for filtering before returning the lymph back to the venous system.

The abdomen and pelvis contain approximately 250 lymph nodes. The lymphatic vessels of the uterus, uterine tubes, and ovaries drain to the external and internal iliac nodes, obturator nodes, para-aortic nodes, and superficial inguinal nodes.<sup>12</sup>

The suspensory ligaments of the urogenital system are important in the mobility and function of the pelvic organs. The uterovesical ligaments attach the bladder to the uterus. The uterosacral ligaments help to suspend the uterus posteriorly. The urogenital system is also supported by ovarian ligaments, suspensory ligaments, and tubo-ovarian ligaments. Just as the ligaments are important in the structure and function of a joint, they are equally important in the mobility and function of the pelvic organs.<sup>14</sup>

The symptoms of dysfunction in the reproductive system can present within the body as dysfunction of the reproductive organs, pelvic asymmetry, sacral dysfunction, bloating, or pain.<sup>14,15</sup> Symptoms related to lymphatic congestion in the pelvic region with hormonal bias are dysmenorrhea, premenstrual syndrome, ovarian cysts, emotional instability, and depression.<sup>16(p795)</sup> Release of fascial and ligamentous restrictions can decrease pressure on blood vessels, thereby optimizing the vascular phase and improving the efficacy of the lymphatic system.<sup>16(p797)</sup> This improved efficacy, in turn, aids in restoring optimal blood flow to the organs, normalizing the ability for hormone production.<sup>15</sup>

Decongestion of the lymphatic system can help remove waste from the organs and thus help normalize their function.<sup>11</sup> Mobilizing fluid and cellular waste from the pelvic cavity should also

allow hormones to more efficiently arrive at the target tissues.<sup>11</sup> Within the reproductive system, this decongestion could theoretically lead to normalized hormone levels, normalized menstrual cycles, and pregnancy. Visceral manipulation, muscle energy, craniosacral therapy, and lymphatic drainage have not been investigated as options for the treatment of infertility; however, the treatment effects of these therapies on other parts of the body could be extrapolated to the pelvis and reproductive system. In the present prospective case series report, I describe the outcomes of 10 infertile women who were treated with manual therapy to the pelvic region.

## **Report of Cases**

### ***Patient Histories and Review of Systems***

Ten infertile women who sequentially presented to a clinic were treated with a standard manual therapy protocol. The women were considered infertile according to the CDC's definition (ie, unable to conceive after 1 year of unprotected intercourse), were not undergoing any other therapies, and had partners who had been tested and found to have normal sperm counts. None of the women had reversals of tubal ligations. There were no known reproductive abnormalities. Any infertility testing was done independently by the women's personal physicians. Of the 10 women treated, 7 had previous pregnancies. Of those 7 who had previously become pregnant, 5 had miscarriages. Five of the women had undergone unsuccessful infertility treatments in the past. At the time of treatment, none of the women were undergoing treatment with hormones, IUI, or IVF. When asked about their past medical history, 5 women listed low back pain, 1 listed irregular periods, 1 listed ovarian tumors, and 1 listed stage IV endometriosis. Two women listed no issues in their medical histories. Regarding surgical history, 1 woman had a dilation and curettage, and 1 woman had a previous cesarean section. A breakdown of the women's past medical and surgical histories is presented in *Table 1*.

**Table 1.**  
**Past Medical and Surgical Histories of Infertile Women Who Received Manual Therapy**

<b>Patient No.</b>	<b>Past Medical History</b>	<b>Past Surgical History</b>	<b>Time Attempting Conception</b>	<b>Type of Infertility</b>	<b>Maternal Age, y</b>
1	Unremarkable	None	1 y	Primary	30
2	Low back pain	Cesarean section	1 y	Secondary	34
3	Right ovarian Tumor, low back pain, miscarriage	Dilation and curettage, laparoscopy (right ovarian tumor removed)	2 y	Secondary	41
4	Irregular cycle, miscarriage	Electrophysiology study ablation, breast augmentation	6 y	Secondary	28
5	Miscarriage	None	3 y	Secondary	36
6	Endometriosis	Pelvic laparoscopy	2 y, 9 mo	Primary	33
7	Miscarriage	None	1 y	Secondary	34
8	Miscarriage	None	7 y	Secondary	39
9	Unremarkable	None	1 y	Secondary	36
10	Low back pain	None	3 y	Primary	41

### ***Evaluation and Intervention***

Women were evaluated for pelvic symmetry at the anterior superior iliac spine and posterior superior iliac spine, osseous restrictions at the sacroiliac joints by means of sacral mobility, visceral fascial restrictions, myofascial trigger points around the entire pelvis, and lymphatic congestion of the uterus and inguinal, iliac, and para-aortic nodes. Manual therapy techniques were chosen according to the specific findings of somatic dysfunction. The techniques were performed according to the protocols described by Chikly,<sup>11</sup> Barral,<sup>15</sup> Upledger and Vredevoogd,<sup>17</sup> and D'Ambrogio.<sup>18</sup>

To attempt increased specificity with evaluation, the mobility of the viscera was recorded as normal, minimally restricted, moderately restricted, or severely restricted. Viscera that was minimally restricted had less than 25% of its motion limited, viscera that was moderately restricted had 25% to 75% of motion restricted, and viscera that was severely restricted was had greater than 75% of mobility restricted.

I evaluated and treated each woman according to the following protocol:

1. Assessed the pelvis for asymmetry and corrected asymmetry with muscle energy techniques, if needed
2. Assessed sacral mobility and corrected dysfunction with craniosacral techniques,<sup>17</sup> if needed
3. Assessed for trigger points around and within the pelvis and treated trigger points with positional release techniques,<sup>18</sup> if needed
4. Assessed lymph drainage of the pelvis and pelvic organs and applied manual lymph drainage techniques,<sup>11</sup> if needed
5. Assessed mobility and motility of pelvic viscera and used fascial techniques to release restrictions,<sup>14</sup> if needed
6. Reassessed symmetry and all mobilities
7. Treatments were repeated twice per week until evaluation revealed unrestricted mobility, symmetry in the pelvis, and normal lymph flow.

The patients were followed up for 3 months after treatments concluded. Women who became pregnant within 3 months were followed up after delivery.

### ***Outcomes***

On examination, 7 of the women were found to have sacral restrictions ranging from mild to severe. Seven women also had restriction in uterine mobility, which ranged from mild to severe. When the lymphatic flow was assessed in the uterus and the pelvis, 1 woman was found to have uterine lymphatic congestion, 1 had mild pelvic lymphatic congestion, 1 had moderate pelvic lymphatic congestion, and 2 had severe pelvic lymphatic congestion. One woman received 1 treatment session, 3 received 2 sessions, 4 received 4 sessions, and 2 received 6 sessions. Six of the 10 women

conceived, had singleton pregnancies, and delivered at full term. Examination findings, number of treatment sessions, and patient outcomes are presented in *Table 2*.

**Table 2.**  
**Profile and Physical Findings of Infertile Women Who Received Manual Therapy**

Full Patient No.	Osseous Issues	Visceral Fascia	Lymphatics	No. of		Delivered
				Treatments	Conceived <sup>a</sup>	
1	Moderate sacral restriction	Normal Mobility	Normal	2	Y	Y
2	Moderate sacral restriction	Mild decreased uterine mobility	Mild pelvic congestion	4	Y	Y
3	Moderate sacral restriction	Mild decreased uterine mobility	Normal	4	N	NA
4	Moderate sacral restriction	Severe decreased uterine mobility	Severe pelvic congestion	4	N	NA
5	Mild sacral restriction	Severe decreased uterine mobility	Severe pelvic congestion	4	N	NA
6	Severe sacral restriction	Severe decreased uterine mobility	Uterus congestion	6	N	NA
7	Normal	Normal mobility	Moderate pelvic congestion	2	Y	Y
8	Normal	Mild decreased uterine mobility	Moderate pelvic congestion	2	Y	Y
9	Normal	Mild decreased uterine mobility	Normal	1	Y	Y
10	Severe sacral restriction	Severe decreased uterine mobility	Normal	6	Y	Y

<sup>a</sup> Conceived within 3 months of last manual therapy session.

**Abbreviation:** NA, not applicable.

### Comment

Although research exists for acupuncture as a treatment for women with infertility,<sup>19,20,21,22,23</sup> little research has been published on the use of manual therapy techniques applied to the pelvis as a

therapeutic treatment option for infertility. A literature search yielded a single published study<sup>24</sup> on physical therapy as a treatment for infertility. In the study, treatment techniques were not disclosed.

In the present case series' population, 6 of 10 previously infertile women were able to conceive within 3 months after receiving various manual therapy techniques to the pelvis. This fertility rate of 60% within a 3-month period is the same as that of fertile couples.<sup>4</sup> These findings suggest that manual therapy applied to the pelvis could be a viable treatment option for infertile women and should be investigated further.

Of note, the manual therapy protocol used in the present case series was limited to external treatments. Other mobilizations not used in the treatment of these women may also be of benefit. For example, internal mobilization of the cervix could allow for better movement of the sperm through the uterus during and after coitus. Arterial mobilizations described by osteopathic physicians Barral and Croibier<sup>25</sup> may also help increase vascularization of the reproductive organs.

One limitation of the assessment used in the present case series is the quantification of the mobility of the organs. No publication could be found referencing a manner to measure visceral mobility. However, Stone<sup>26</sup> states that testing of visceral mobility involves trying to replicate “normal” movement and trying to assess whether the motion appears to be restricted or altered. This assessment allows for a before-and-after comparison of the tissues. She also states, “Altered physical properties arising from inflammation and other pathological processes will affect palpable visceral characteristics such as stretch, deformation and compressibility. In this way, osteopaths can distinguish to some degree whether a tissue is normal or not normal.”<sup>26</sup> Further research would be beneficial in assessing range of motion of organs to obtain normative data.

Another limitation of this study, as with any case series report, is the low number of women treated. It is difficult to say with a high level of certainty that the manual therapy techniques were what contributed to the improved fertility. Studies with larger groups, including control participants,

would help to discern the exact statistical significance of manual techniques in the management of infertility.

In the present study, the number of treatments for each woman ranged from 1 to 6. This low number of treatments suggests that manual medicine would be an inexpensive treatment option for infertility. In addition, manual mobilization techniques do not contain the risks associated with traditional infertility treatments such as multiple births.

Just as manual therapy is used as a first course of treatment for patients considering back surgery,<sup>27</sup> manual therapy could potentially be used as a first course of treatment for infertile patients considering hormone therapy, IUI, or IVF treatments. The ethics committee for the American Society for Reproductive Medicine states, “For those treatments with very poor success rates, clinicians must be vigilant in their presentation of risks, benefits, and alternatives.”<sup>28</sup> If manual medicine is shown to be an effective yet inexpensive means to increase fertility rates in the infertile population, then it could be considered as a primary treatment option for infertile women, rather than as an alternative treatment option.

## **Conclusion**

For 6 of the 10 women in the present case series report, fertility rates improved after manual therapy was applied to the pelvic area. Further research is needed to assess the efficacy of manual therapy as a treatment option for infertile women. For future studies, researchers should use a larger study population, include a control group, and assess the efficacy of the individual manual techniques.

**Financial Disclosures:** None reported.

Address correspondence to Mary Ellen Kramp, DPT, Certified Lymphedema Therapist-Lymphology Association of North America, Relief at Hand, LLC, 8215 113th St, Seminole, FL 33772-4128.

E-mail: maryellenpt@aol.com

## References

1. Centers for Disease Control and Prevention. Infertility. Centers for Disease Control and Prevention Web site. Updated July 18, 2012. <http://www.cdc.gov/nchs/fastats/fertile.htm>. Accessed August 16, 2012.
2. Dey SK. How we are born. *J Clin Invest.* 2010;120(4):952-955.
3. Katz V, Lentz GM, Gershenson D, Lobo R. *Comprehensive Gynecology.* 5th ed. Philadelphia, PA: Mosby Elsevier; 2007.
4. Evaluation of the infertile couple. In: Schorge JO, Schaffer JI, Halvorson LM, Hoffman BL, Bradshaw KD, Cunningham FG. *Williams Gynecology.* New York, NY: McGraw Hill; 2008:426.
5. Collins JA, Burrows EA, Wilan AR. The prognosis for live birth among untreated infertile couples. *Fertil Steril.* 1995;64(1):22-28.
6. van der Steeg JW, Steures P, Eijkemans MJ, et al. Pregnancy is predictable: a large-scale prospective external validation of the prediction of spontaneous pregnancy in subfertile couples. *Hum Reprod.* 2007;22(2):536-542.
7. Katz P, Showstack J, Smith JF, et al. Costs of infertility treatment: results from an 18-month prospective cohort study [published online ahead of print December 4, 2010]. *Fertil Steril.* 2011;95(3):915-921..
8. Baker VL, Jones CE, Cometti B, et al. Factors affecting success rates in two concurrent clinical IVF trials: an examination of potential explanations for the difference in pregnancy rates between the United States and Europe. *Fertil Steril.* 2010;94(4):1287-1291.
9. Centers for Disease Control and Prevention. Assisted Reproductive Technology report: national summary report. Centers for Disease Control and Prevention Web site. <http://apps.nccd.cdc.gov/art/Apps/NationalSummaryReport.aspx>. Accessed September 19, 2012.

10. Centers for Disease Control and Prevention. Assisted Reproductive Technology report, section 2: ART cycles using fresh, nondonor eggs or embryos (part A). Centers for Disease Control and Prevention Web site. [http://www.cdc.gov/art/ART2009/sect2\\_fig6-15.htm#10](http://www.cdc.gov/art/ART2009/sect2_fig6-15.htm#10). Accessed September 19, 2012.
11. Chikly B. *Silent Waves: Theory and Practice of Lymph Drainage Therapy, an Osteopathic Lymphatic Technique*. Scottsdale, AZ: International Health & Healing Inc; 2005.
12. Borley NR, Healy JC. Female reproductive system. In: Standring S, ed. *Gray's Anatomy: The Anatomical Basis of Clinical Practice*. 40th ed. London, England: Churchill Livingstone Elsevier; 2008:1279-1304.
13. Zkink JG. Applications of the osteopathic holistic approach to homeostasis. *Am Aca of Osteo Yearb*. 1973;37-47.
14. Barral JP. *Visceral Manipulation II*. Seattle, WA: Eastland Press; 1989.
15. Barral JP. *Urogenital Manipulation*. Seattle, WA: Eastland Press; 1993.
16. Chila AG, executive ed. *Foundations of Osteopathic Medicine. 3rd ed*. Baltimore, MD: Lippincott Williams & Wilkins; 2011.
17. The spinal dura matter and sacrococcygeal complex. In: Upledger JE, Vredevoogd JD. *Craniosacral Therapy*. Seattle, WA: Eastland Press; 1996:131-151.
18. D'Ambrogio KJ, Roth GB. *Positional Release Therapy*. St. Louis, MO: Mosby; 1997.
19. Beal MW. Women's use of complementary and alternative therapies in reproductive health care. *J Nurse Midwifery*. 1998;43(3):224-234.

20. Birkeflet O, Laake P, Vøllestad N. Traditional Chinese medicine patterns and recommended acupuncture points in infertile and fertile women. *Acupunct Med.* 2012;30(1):12-16.
21. Cochrane S, Smith CA, Possamai-Inesedy A. Development of a fertility acupuncture protocol: defining an acupuncture treatment protocol to support and treat women experiencing conception delays. *J Altern Complement Med.* 2011;17(4):329-337.
22. Franconi G, Manni L, Aloe L, et al. Acupuncture in clinical and experimental reproductive medicine: a review [published online ahead of print February 4, 2011]. *J Endocrinol Invest.* 2011;34(4):307-311.
23. Bovey M, Lorenc A, Robinson N. Extent of acupuncture practice for infertility in the United Kingdom: experiences and perceptions of the practitioners [published online ahead of print May 10, 2010]. *Fertil Steril.* 2010;94(7):2569-2573.
24. Wurn BF, Wurn LJ, King R, et al. Treating female infertility and improving IVF pregnancy rates with a manual physical therapy technique. *MedGenMed.* 2004;6(2):51.
25. Barral JP, Croibier A. *Visceral Vascular Manipulations.* New York, NY: Churchill Livingstone Elsevier; 2011.
26. General principles. In: Stone CA. *Visceral and Obstetric Osteopathy.* New York, NY: Elsevier; 2007:20.
27. Issack PS, Cunningham ME, Pumberger M, Hughes AP, Cammisa FP Jr. Degenerative lumbar spinal stenosis: evaluation and management. *J Am Acad Orthop Surg.* 2012;20(8):527-535.
28. Ethics Committee of the American Society for Reproductive Medicine. Fertility treatment when the prognosis is very poor or futile. *Fertil Steril.* 2004;82(4):806-810.

### **CME Questions**

1. Worldwide, what percentage of couples are infertile?
  - (a) 5%
  - (b) 22%
  - (c) 15%**

(d) 33%

2. What percentage of fertile couples will conceive within a 3-month period?

(a) 33%

**(b) 57%**

(c) 72%

(d) 20%